

Note on multiple additivity of minimal Renyi entropy output of the Werner–Holevo channels

R. Alicki¹ and M. Fannes²

*Institute for Theoretical Physics and Astrophysics, University of Gdańsk,
ul. Wita Stwosza 57, PL-80-952 Gdańsk, Poland*

*Instituut voor Theoretische Fysica, K.U.Leuven, Celestijnenlaan 200D,
B-3001 Heverlee, Belgium*

Abstract: We give an elementary self-contained proof that the minimal entropy output of arbitrary products of channels $\rho \mapsto \frac{1}{d-1}(1-\rho^T)$ is additive.

This paper is concerned with efficiency of transmission of classical information through a particular quantum channel. Generally, the minimal entropy output of a general quantum channel Γ , given in terms of a completely positive trace-preserving map on the d -dimensional density matrices, quantifies the minimal noise produced by the channel in terms of the von Neumann entropy of transformed pure states

$$\text{MEO}[\Gamma] := \inf_{\varphi, \|\varphi\|=1} S(\Gamma(|\varphi\rangle\langle\varphi|))$$

where $S(\rho)$ is the von Neumann entropy of a density matrix ρ . Recalling the definition of the p -Renyi entropy of a density matrix ρ , $p > 1$

$$S_p(\rho) := -\frac{1}{p-1} \log(\text{Tr } \rho^p),$$

the minimal p -Renyi entropy output is

$$\text{MEO}_p[\Gamma] := \inf_{\varphi, \|\varphi\|=1} S_p(\Gamma(|\varphi\rangle\langle\varphi|)). \quad (1)$$

Obviously, $\lim_{p \downarrow 1} \text{MEO}_p[\Gamma] = \text{MEO}[\Gamma]$ and we can therefore extend the notation MEO_p to $p = 1$.

The p -Renyi entropy of a density matrix is directly related to its p -norm. Let X be a general, not necessarily square, complex matrix. The p -norm of X is given by

$$\|X\|_p := \left(\text{Tr} |X|^p \right)^{\frac{1}{p}} = \left(\sum_j \chi_j^p \right)^{\frac{1}{p}}$$

¹E-mail: fizra@univ.gda.pl

²E-mail: mark.fannes@fys.kuleuven.ac.be

where $|X| = \sqrt{X^*X}$ and where the χ_j are the (non-zero) singular values of X . The p -norm is the non-commutative analogon of the usual p -norm of sequences of complex numbers. As a density matrix is positive semi-definite, we have the immediate relation

$$S_p(\rho) = -\frac{p}{p-1} \log \|\rho\|_p. \quad (2)$$

In terms of p -norms, the quantity corresponding to $\text{MEO}_p(\Gamma)$ is

$$\nu_p(\Gamma) := \sup_{\varphi, \|\varphi\|=1} \left\| \Gamma(|\varphi\rangle\langle\varphi|) \right\|_p.$$

Using (2) and (1) we find

$$\text{MEO}_p(\Gamma) = -\frac{p}{p-1} \log \nu_p(\Gamma). \quad (3)$$

The additivity conjecture

$$\text{MEO}[\Gamma_1 \otimes \Gamma_2] = \text{MEO}[\Gamma_1] + \text{MEO}[\Gamma_2] \quad (4)$$

means that entanglement cannot help to lower the noise in parallel channels, described by tensor products. Recently, the relevance of this conjecture has been stressed, as it was shown to be equivalent to other important additivity issues in quantum information theory, such as Holevo capacity and entanglement of formation, see [6]. There is a similar conjecture for MEO_p , $1 < p \leq 2$. As both the von Neumann and the Renyi entropies are additive on tensor products, the conjecture amounts to showing that

$$\text{MEO}_p[\Gamma_1 \otimes \Gamma_2] \geq \text{MEO}_p[\Gamma_1] + \text{MEO}_p[\Gamma_2]. \quad (5)$$

The relation (3) allows us to restate the conjecture (4) as multiplicativity of maximal p -norms

$$\nu_p[\Gamma_1 \otimes \Gamma_2] = \nu_p[\Gamma_1] \nu_p[\Gamma_2].$$

We shall consider here a particular quantum channel, known as Werner–Holevo channel

$$\Gamma_d(\rho) = \frac{1}{d-1} (\mathbb{1} - \rho^\top). \quad (6)$$

Here, $A^\top$ denotes the transpose of A w.r.t. the standard basis of $\mathbb{C}^d$. The map Γ_d has both a high symmetry and a truly entangling nature compared e.g. with the depolarising channel. In fact, $U\Gamma_d(\rho)U^* = \Gamma_d(\overline{U}\rho(\overline{U})^*)$ where

U is an element of $SU(d)$ and $\bar{U}$ is its complex conjugate and it is the non-trivial extreme point in the set of all such maps. In the case $d = 3$, it is even an extreme point in the set of all completely positive trace-preserving maps. This channel has often been considered in the literature, see e.g. [2, 7, 1]. We show that the minimal entropy output of $\otimes^N \Gamma_{d_j}$ is multiply additive for $1 \leq p \leq 2$

$$\text{MEO}_p[\otimes_{j=1}^N \Gamma_{d_j}] = \sum_{j=1}^N \text{MEO}_p[\Gamma_{d_j}]. \quad (7)$$

In the particular case $p = 1$, $N = 2$ and $d_1 = d_2$, formula (7) has been proven in [1] using more advanced techniques, see also [4]. Because of (3) this is equivalent to showing the multiple multiplicativity of the maximal p -norms, see [3]

$$\nu_p[\otimes_{j=1}^N \Gamma_{d_j}] = \prod_{j=1}^N \nu_p[\Gamma_{d_j}], \quad 1 < p \leq 2.$$

A straightforward computation yields

$$\begin{aligned} \text{MEO}_p[\Gamma_d] &:= \inf_{\varphi, \|\varphi\|=1} S_p\left(\Gamma_d(|\varphi\rangle\langle\varphi|)\right) \\ &= \inf_{\varphi, \|\varphi\|=1} S_p\left(\frac{1}{d-1}(\mathbb{1} - |\bar{\varphi}\rangle\langle\bar{\varphi}|)\right) = \log(d-1). \end{aligned}$$

Here, $\bar{\varphi}$ is the complex conjugate of φ in the standard basis.

Next, we write out the action of the channel $\otimes_{j=1}^N \Gamma_{d_j}$ on the pure state generated by a normalised vector $\Omega \in \otimes_{j=1}^N \mathbb{C}^{d_j}$

$$X_N(\Omega) := \otimes_{j=1}^N \Gamma_{d_j}(|\Omega\rangle\langle\Omega|) = \prod_{j=1}^N \frac{1}{d_j-1} \sum_{\Lambda \subset \{1,2,\dots,N\}} (-1)^{|\Lambda|} \rho_\Lambda \otimes \mathbb{1}_{\Lambda^c}.$$

Here, ρ_Λ is the reduced density matrix of $|\bar{\Omega}\rangle\langle\bar{\Omega}|$ to the sites in Λ . We now compute

$$\begin{aligned} &\text{Tr}(X_N(\Omega))^2 \\ &= \prod_{j=1}^N \frac{1}{(d_j-1)^2} \sum_{\Lambda, \Lambda' \subset \{1,2,\dots,N\}} (-1)^{|\Lambda|+|\Lambda'|} \text{Tr}\left((\rho_\Lambda \otimes \mathbb{1}_{\Lambda^c})(\rho_{\Lambda'} \otimes \mathbb{1}_{\Lambda'^c})\right) \\ &= \prod_{j=1}^N \frac{1}{(d_j-1)^2} \sum_{\Lambda, \Lambda' \subset \{1,2,\dots,N\}} (-1)^{|\Lambda|+|\Lambda'|} \left(\prod_{j \in (\Lambda \cup \Lambda')^c} d_j\right) \text{Tr} \rho_{\Lambda \cap \Lambda'}^2 \end{aligned}$$

$$\begin{aligned}
&= \prod_{j=1}^N \frac{1}{(d_j - 1)^2} \sum_{\Lambda \subset \{1, 2, \dots, N\}} \text{Tr} \rho_\Lambda^2 \sum_{\Delta \subset \Lambda^C} (-1)^{|\Delta|} \sum_{\Delta' \subset \Lambda^C \setminus \Delta} (-1)^{|\Delta'|} \prod_{j \in (\Lambda^C \setminus \Delta) \setminus \Delta'} d_j \\
&= \prod_{j=1}^N \frac{1}{(d_j - 1)^2} \sum_{\Lambda \subset \{1, 2, \dots, N\}} \text{Tr} \rho_\Lambda^2 \prod_{j \in \Lambda^C} (d_j - 2) \leq \prod_{j=1}^N \frac{1}{d_j - 1}. \tag{8}
\end{aligned}$$

In this computation Λ^C denotes the complement of Λ in $\{1, 2, \dots, N\}$ and we used $\text{Tr}(\rho_\Lambda)^2 \leq 1$.

Finally, we use the inequality

$$S_2(\rho) \leq S_p(\rho) \leq S(\rho), \quad 1 \leq p \leq 2$$

relating the p -Renyi entropies and the 2-Renyi entropy [5] to obtain

$$\text{MEO}_p[\otimes_{j=1}^N \Gamma_{d_j}] \geq -\log \left(\prod_{j=1}^N \frac{1}{d_j - 1} \right) = \sum_{j=1}^N \text{MEO}_p[\Gamma_{d_j}],$$

proving hereby (7).

Acknowledgements: R.A. is grateful for the hospitality during his stay at the K.U.Leuven and acknowledges the financial support by the grant PBZ-MIN-008/P03/2003 of the Polish Ministry of Scientific Research and Information Technology and EC grant RESQ contract IST-2001-37559. This work was also partially supported by F.W.O., Vlaanderen grant G.0109.01. The authors also thank the referee for helpful suggestions concerning the presentation of their result.

References

- [1] N. Datta, A.S. Holevo and Y.M. Suhov: A quantum channel with additive minimum output entropy, quant-ph/0403072
- [2] A.S. Holevo: Remarks on the classical capacity of quantum channel, quant-ph/0212025
- [3] C. King and M.B. Ruskai: Comments on multiplicativity of maximal p -norms when $p = 2$, quant-ph/0401026
- [4] K. Matsumoto T. Shimono and A. Winter: Remarks on additivity of the Holevo channel capacity and of the entanglement of formation, *Commun. Math. Phys.* **246**, 427–442 (2004)

- [5] A. Renyi: *Probability Theory*, Akademiai Kiado, Budapest, North-Holland, Amsterdam (1970)
- [6] P. Shor: Equivalence of additivity questions in quantum information theory, quant-ph/0305035
- [7] R.F. Werner and A.S. Holevo: Counterexample to an additivity conjecture for output purity of quantum channels, *J. Math. Phys.* **43**, 4353 (2002) quant-ph/0203003